PRESCRIPTION AND CERTIFICATION OF MEDICAL NECESSITY ORDER

Patient Name:	Rep:	_ Doctor Name:
Surgery / Procedure:	Surgery Date:	NPI #:
OOB:	Patient Diagnosis:	
	ICDx 10 Code #:	
Home Intermittent Limb Compression		anaFlexx TENS / NMES Device (E0745)
Other	Ot	her
Caprini Risk Factor Scoring - Please ched	ck each relevant Risk Factor Score for	this patient
EACH RISK FACTOR = 1 POINT	EACH RISK FACTOR = 2 POINTS	EACH RISK FACTOR = 3 POINTS
Age 40 − 60 Years	O Age 61 – 69 Years	O Age 70 + Years
Obesity (BMI > 25)	○ BMI > 40	○ BMI > 50
Use of Tobacco / Smoker	☐ Immobility Ambulate <100ft 3 times per day	Malignancy and / or Chemotherapy
Anesthesia Time < 1 Hour	○ Major Surgery > 60 Minutes	○ Sepsis
Leg Swelling, Ulcers, Varicose Veins	Anesthesia Time > 30 Minutes	 Congestive Heart Failure or Myocardial Infarction
Pregnancy or Postpartum < 1 Month	Acute Respiratory Failure / Severe COPD	Hypercoagulable Syndrome
Estrogen Therapy / Birth Control Pill / HRT	Arthroscopic Surgery > 60 minutes	Family or Personal History VTE
Nephrotic Syndrome (> 4GM Proteinuria/Day)	○ Laparoscopic Surgery > 60 Minutes	
Insufflation of Abdomen During Laparoscopic Surge	Y O Anticipated Bed Confinement / Immobili	zation > 24Hrs
Acute Infection other than Sepsis		
Inflammatory Bowel Disease	EACH RISK FACTOR = 5 POINTS	HIGH RISK
Use of Central Venous Catheter	Stroke with Paralysis	= 3 + Points 🖳
Use of Tourniquet	Major Trauma or Spinal Cord Injury	
Dehydration	Hip, Pelvis, or Leg Fracture	TOTAL RISK POINTS
Leg Plaster cast or brace	Knee or Hip Arthroplasty	Low Risk 1 Point or Less
Minor Surgery	Major Surgery > 3 Hours	Medium Risk 2 Points
Patient on Current Bed Rest	OTHER RISK FACTORS = 1 POINT	○ High Risk 3-4 Points
		○ Very High Risk 5 + Points
	O	
 Reduce swelling and pain. Fewer transfusions of allogenic blood The patient will continue routine post The patient will use the device as instantimized. 	nent indicates the use of mechanical thronounces. I am prescribing this device to maxinollowing major orthopedic/neurological sund DVT management following surgery. d. coperative follow-up with me for clinical extructed, daily at home, for 3-months or unto prevention are contraindicated following mechanical necessary for home use for the above in	nboprophylaxis by pneumatic device and mize the outcome of surgery and minimize urgery: valuation. til adequately mobilizing and risk of DVT/PE is
Doctor's Name :	NPI#: En	nail:

I hereby authorize the release af patient health information from my office to Monarch Medical Solutions for the purposes of filing claims with patient's insurance company. This information will be shared for the time it takes to file and/or appeal my claim. I understand that I have the right to revoke this authorization, in writing anytime, except where uses or disclosures have already been made based an my original permission.